

Data collection form – Patients referred for a spacer device, PAAP and/or has been prescribed three or more short-acting bronchodilators without any corticosteroid within a six-month period

Patient bag label	Date of intervention	Date of referral	Reason for referral	GP practice	Action taken following the intervention, e.g. inhaler technique check, NMS
			Has not been prescribed a spacer device for use with a press and breathe pressurised MDI (the patient is aged 5-15 years) <input type="checkbox"/> Does not have a Personalised Asthma Action Plan <input type="checkbox"/> Has been prescribed three or more short-acting bronchodilator inhalers without any corticosteroid inhaler within a six-month period <input type="checkbox"/>		
			Has not been prescribed a spacer device for use with a press and breathe pressurised MDI (the patient is aged 5-15 years) <input type="checkbox"/> Does not have a Personalised Asthma Action Plan <input type="checkbox"/> Has been prescribed three or more short-acting bronchodilator inhalers without any corticosteroid inhaler within a six-month period <input type="checkbox"/>		
			Has not been prescribed a spacer device for use with a press and breathe pressurised MDI (the patient is aged 5-15 years) <input type="checkbox"/> Does not have a Personalised Asthma Action Plan <input type="checkbox"/> Has been prescribed three or more short-acting bronchodilator inhalers without any corticosteroid inhaler within a six-month period <input type="checkbox"/>		
			Has not been prescribed a spacer device for use with a press and breathe pressurised MDI (the patient is aged 5-15 years) <input type="checkbox"/> Does not have a Personalised Asthma Action Plan <input type="checkbox"/> Has been prescribed three or more short-acting bronchodilator inhalers without any corticosteroid inhaler within a six-month period <input type="checkbox"/>		
			Totals		